

Longitudinal assessment of the impact of orthodontic treatment on adolescents' quality of life: a comparison between boys and girls using a condition specific questionnaire

Larissa Corradi-Dias¹, Saul M Paiva¹, Alexandre F Drummond², José Alcides A de Arruda³, Leonardo FS de Menezes², Lucas G Abreu¹

1. Universidade Federal de Minas Gerais, Faculdade de Odontologia, Departamento de Saúde Bucal da Criança e do Adolescente, Belo Horizonte, Brasil

2. Universidade Federal de Minas Gerais, Faculdade de Odontologia, Departamento de Odontologia Restauradora, Belo Horizonte, Brasil

3. Universidade Federal de Minas Gerais, Faculdade de Odontologia, Departamento de Clínica, Patologia e Cirurgia Odontológicas, Belo Horizonte, Brasil

ABSTRACT

The aim of this study was to compare the impact of the first year of wearing of a fixed orthodontic appliance on the Oral Health-Related Quality of Life (OHRQoL) between boys and girls, by means of a condition-specific instrument. The study included 69 adolescents aged 10 to 18 years, who were undergoing orthodontic treatment with a fixed appliance. Of the 69 adolescents, 38 were girls (55.1%) and 31 were boys (44.9%). They answered the Brazilian version of the Impact of Fixed Appliance Measure (B-IFAM) questionnaire three months (T1) and one year (T2) after the fixed appliance was installed. This questionnaire contains 43 questions, distributed across nine domains. The higher the scores, the more negative the perception of the adolescent concerning the impact of the fixed appliance on his/her OHRQoL. Sociodemographic and clinical variables were also analyzed, and statistical analysis was performed. For the domains, the effect size (the magnitude of the difference between girls and boys) and the minimal clinically important difference were also calculated. The adjusted regression showed that there was a significantly greater increase in the overall B-IFAM score in girls than in boys, indicating a more negative perception of the OHRQoL over the study time [Coefficient=11.77 (3.47–20.60), $p=0.006$]. From T1 to T2, there was a significantly greater increase in the scores (more negative perception of OHRQoL over time) in girls than in boys for the domains aesthetics ($p=0.034$) and physical impact ($p=0.011$). These differences were clinically significant. The effect size (the magnitude of the difference) was moderate. The impact of wearing a fixed appliance on the OHRQoL was more negative in girls than in boys during the first year of orthodontic treatment.

Keywords: adolescent - quality of life - orthodontic treatment - fixed appliance

To cite:

Corradi-Dias L, Paiva SM, Drummond AF, de Arruda JAA, de Menezes LFS, Abreu LG. Longitudinal assessment of the impact of orthodontic treatment on adolescents' quality of life: a comparison between boys and girls using a condition specific questionnaire. Acta Odontol Latinoam. 2022 Sep 30;35(2):125-133. <https://doi.org/10.54589/aol.35/2/125>

Corresponding Author:

Lucas G. Abreu
lucasgabreu01@gmail.com

Received: April 2022.

Accepted: June 2022.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License

Avaliação longitudinal do impacto do tratamento ortodôntico na qualidade de vida de adolescentes: comparação entre meninos e meninas utilizando um questionário condição específica

RESUMO

O objetivo deste estudo foi comparar o impacto do primeiro ano de uso do aparelho fixo na qualidade de vida relacionada a saúde bucal (QVRSB) entre meninas e meninos, através de um instrumento condição específica. Sessenta e nove adolescentes entre 10 e 18 anos, em tratamento ortodôntico com aparelho fixo foram incluídos. Adolescentes responderam ao questionário Impact of fixed appliance measure (B-IFAM) no terceiro mês de uso do aparelho fixo (T1) e um ano após a colagem do aparelho fixo (T2). Este questionário possui 43 perguntas, distribuídas em nove domínios. Quanto maior os escores, mais negativa a percepção do adolescente com relação ao impacto do aparelho fixo na QVRSB. Variáveis sociodemográficas e clínicas também foram avaliadas. Análise estatística foi realizada. Para os domínios, tamanho de efeito (a magnitude da diferença entre meninas e meninos) e diferença mínima clinicamente importante também foram calculadas. Dos 69 adolescentes, 38 eram meninas (55,1%) e 31 eram meninos (44,9%). Na regressão ajustada, meninas apresentaram um aumento significativamente maior do escore total do B-IFAM do que meninos, indicando uma percepção mais negativa da QVRSB ao longo do tempo de acompanhamento [Coeficiente=11,77 (3,47–20,60), $p=0.006$]. Meninas apresentavam um aumento significativamente maior dos escores de T1 para T2 (percepção mais negativa da QVRSB ao longo do tempo) em relação aos meninos para os domínios estética ($p=0,034$) e impacto físico ($p=0,011$). Estas diferenças também foram clinicamente significativas. O tamanho do efeito (a magnitude da diferença) foi moderado. Meninas demonstraram um impacto mais negativo do uso do aparelho fixo na QVRSB do que meninos nos 12 primeiros meses de tratamento ortodôntico.

Palavras-Chave: adolescente, qualidade de vida - tratamento ortodôntico - aparelho fixo.

INTRODUCTION

There is increasing scientific interest in people's wellbeing, and research seeks to understand the relationship between multidimensional constructs such as quality of life or self-esteem, and health¹. Oral health conditions have been found to exert major impact on quality of life, giving rise to the term oral health-related quality of life (OHRQoL)^{2,3}. The literature has shown the negative impact of oral conditions such as malocclusion on the OHRQoL of young people, with strong repercussions on oral functions, and emotional and social wellbeing^{4,5}. Orthodontic treatment with a fixed appliance seeks to correct tooth positions and skeletal discrepancies, providing more functionally and esthetically favorable occlusion, and, consequently, better quality of life⁶.

Changes in OHRQoL have been observed during the orthodontic treatment of adolescents wearing fixed appliances. Deterioration in quality of life is common during the first months of wearing an orthodontic device, mainly due to the oral symptoms and functional limitations it causes⁷. However, interestingly, other studies have found a positive impact on adolescent emotional wellbeing, even during the initial stages of orthodontic therapy, thanks to their expectation of having their teeth corrected^{7,8}. The longitudinal studies available in the literature evaluate the impact of orthodontic treatment on adolescent quality of life by means of generic questionnaires to evaluate a condition-specific outcome of wearing a fixed appliance. Condition-specific questionnaires are more sensitive and more responsive, and thus provide more reliable results for evaluating a construct as complex as OHRQoL⁹.

A recent cross-sectional study used a condition-specific questionnaire on wearing a fixed appliance to compare the impact of orthodontic treatment between boys and girls in the sixth month of wearing the appliance. The impact on quality of life was more negative in girls than in boys. Girls expressed a more negative perception of the pain caused by the fixed device, difficulty in hygiene, and social impact, especially regarding interaction with their peers¹⁰. In the context of orthodontics, more in-depth evaluations of the differences between boys and girls are relevant, given that they may have different perceptions and expectations regarding health outcomes¹¹. The aim of this longitudinal

study is to compare boys and girls concerning the impact of the first year of orthodontic treatment with a fixed appliance on their OHRQoL, by means of a condition-specific instrument.

METHOD

Study design

This longitudinal study follows the guidelines set forth in *Strengthening the Reporting of Observational Studies in Epidemiology* (STROBE)¹².

Participants, study location, data collection period, and ethics

The sample consisted of 79 male and female adolescents, 10 to 18 years of age, who were in the third month of orthodontic treatment with a fixed appliance at the School of Dentistry of the Federal University of Minas Gerais (UFMG), located in Belo Horizonte, Brazil. Any adolescents with cognitive disorders reported by their parents/guardians or with craniofacial anomalies were excluded from the study. Participants were recruited from January 2017 to December 2018, and data were collected from January 2017 to February 2020. This study was approved by the UFMG Research Ethics Committee concerning research involving human beings (No. 62116216.2.0000.5149).

Study Variables

Dependent Variable: OHRQoL

The impact of wearing a fixed appliance on adolescent quality of life was evaluated with a condition-specific instrument – Impact of Fixed Appliances Measure (IFAM). This instrument was developed in England¹³, translated to Brazilian Portuguese, cross-culturally adapted, and subsequently validated for use in the Brazilian population¹⁴, resulting in a national version of the questionnaire, called B-IFAM.

The B-IFAM consists of 43 questions. The answer choices for each question follow the Likert scale of 1 to 5, with 1 = completely disagree, 2 = disagree, 3 = neither agree nor disagree, 4 = agree, 5 = completely agree. The overall B-IFAM score ranges from 43 to 215. Higher scores indicate more negative perception regarding the impact of wearing a fixed appliance on the respondent's quality of life. These 43 questions can be broken down into nine domains: aesthetics (5 questions), functional limitations (3 questions), dietary impact (6 questions), oral hygiene impact

(3 questions), maintenance (2 questions), physical impact (9 questions), social impact (5 questions), time constraints (5 questions), and travel/cost/inconveniences (5 questions). Each domain can also be scored individually. The B-IFAM was completed by the adolescents, without any help, with the exception of the questions in the two last domains, for which the parents/guardians helped them to answer the questions, as recommended by the authors. The adolescents answered the questionnaire to evaluate the impact of the wearing of a fixed appliance on their quality of life at two distinct times: three months (T1) and 12 months (T2) after beginning orthodontic treatment with a fixed appliance.

Independent variable: sex

Sex was the main independent variable, and participants were divided into two groups: female and male.

Confounding variables: demographic, socioeconomic, and clinical data

In addition to the sex variable, age and family income variables were collected. The family monthly income was evaluated according to the Brazilian minimum salary at the time of data collection, and established by adding together the monthly income of all the active family members. This variable was dichotomized by the median into adolescents whose families had a monthly income ≤ 2 minimum salaries and adolescents whose families had a monthly income > 2 minimum salaries. The age variable was also dichotomized through the median in individuals who were ≤ 12 years old and individuals who were > 12 years old.

During clinical data collection, the adolescents underwent clinical examinations, performed in a room with dental equipment, under artificial light, using a WHO (World Health Organization) probe, and a clinical mirror. During this examination, the orthodontic indication of the extraction of pre-molars (yes/no) and the severity of the malocclusion were evaluated through the Dental Aesthetic Index (DAI)¹⁵. The DAI is a cross-cultural index that enables the evaluation of 10 occlusal characteristics related to dental-facial anomalies, according to three components: dentition (number of absent incisors, canines, and pre-molars), crowding and/or spacing (crowding in the region of the incisors, spacing in the incisor region, diastema between the maxillary central incisors, greater irregularity in the maxillary

anterior teeth, and greater irregularity in the mandibular anterior teeth), as well as occlusion (horizontal trespass, anterior crossbite, open bite, and anterior-posterior molar relationship). Scores for each occlusal characteristic were multiplied by a coefficient and added to the constant of 13 in order to obtain the total score of the DAI for each participant. Based on the DAI scores, the adolescent could be classified into one of four levels of severity of malocclusion: slight malocclusion ($DAI \leq 25$), defined malocclusion ($DAI = 26-30$), severe malocclusion ($DAI = 31-35$), and highly severe malocclusion ($DAI \geq 36$)¹⁵. In this study, the severity of the malocclusion was dichotomized into slight/defined malocclusion ($DAI \leq 30$) and severe/highly severe malocclusion ($DAI \geq 31$). The researcher responsible for data collection underwent a calibration process with another researcher who had experience and ability in the application of DAI. The calibration process took place in two stages, with the first stage being theoretical and the second, clinical. The theoretical stage consisted of discussing the criteria used in the DAI to classify the severity of the malocclusion and the need for orthodontic treatment. The clinical stage consisted of evaluating 15 plaster models and clinical exams of 15 adolescents who were not included in the main study. The evaluations were conducted separately by the two researchers in order to calculate inter-examiner agreement. After one week, the plaster models and the adolescents were re-evaluated by the researcher who collected the data in order to calculate intra-examiner agreement. The Kappa values were 0.80 and 0.90 for inter- and intra-examiner evaluations, respectively, being considered satisfactory values¹⁶. Finally, data on the type of fixed appliance worn (conventional, self-ligating, or aesthetic) were collected.

Pilot Study

A pilot study to evaluate the data collection strategy was conducted with 10 adolescents who were not included in the main study. No change in the data collection strategy was necessary.

Statistical analysis

Statistical analysis was performed using the Statistical Package for Social Science (SPSS, version 25.0, IBM Inc., Armonk, USA). First, the Kolmogorov-Smirnov test demonstrated that

the total B-IFAM scores in T1 and T2, as well as the difference between T1 and T2, had normal distribution. For the latter, the negative sign (-) indicated a more negative perception of the OHRQoL from T1 to T2. Participants with complete data were compared to individuals excluded because of the loss of data, using the Fisher test for categorical variables (sex, age, monthly household income, severity of malocclusion, and orthodontic tooth extraction), and the *t* test for numerical variables (overall B-IFAM score in T1).

Females and males were compared for the variables age, family monthly income, malocclusion severity, and indication of orthodontic extraction of premolars, as well as for the outcome variables overall score of B-IFAM in T1, T2, and the difference between T1 and T2. Fisher's test and Pearson's test were used for categorical variables and the *t* test was used for the numerical variables.

The crude and adjusted associations between sex and the variables difference in the overall B-IFAM score between T1 and T2 were tested with linear regression. Regression coefficients and 95 % confidence intervals (CI) were determined as association measures. The adjusted regression model included the control of the variables age, family monthly income, malocclusion severity,

indication of orthodontic extraction, and the overall B-IFAM score in T1.

The domains of the B-IFAM were also compared between females and males, using the *t* test. The differences between girls and boys, the effect size (the magnitude of the difference), and the 95 % CI were determined. As a reference, effect size values close to 0.20 were small, values close to 0.50 were medium, and values close to 0.80 were large¹⁷. The minimal clinically important difference (MCID) was calculated by multiplying the standard deviation of the domain assessed for the entire sample (pooled standard deviation) by 0.5¹⁸.

Finally, covariance was analyzed for comparisons of the domains of the B-IFAM between females and males, controlling for the type of fixed appliance.

RESULTS

Of the 79 adolescents that began the study, 10 were excluded due to missing data (they did not fill out the B-IFAM at T2, or some other information was not provided). No difference was found between the 69 adolescents who participated in the entire study and the 10 who were excluded, for the variables sex and age, monthly family income, malocclusion severity, indication of tooth extraction, and the overall score of B-IFAM in T1 (Table 1). Of the 69

Table 1. Comparison of adolescents in the study sample to those excluded because of missing data

	Study sample	Excluded	<i>p</i> value
Independent variables	N (%)	N (%)	
Sex			
Girls	38 (55.1)	03 (30.0)	0.183*
Boys	31 (44.9)	07 (70.0)	
Age			
≤12 years	44 (63.8)	05 (55.6)	0.720*
>12 years	25 (36.2)	04 (44.4)	
Family Monthly income			
≤2 minimum wages	44 (63.8)	05 (55.6)	0.720*
>2 minimum wages	25 (36.2)	04 (44.4)	
Malocclusion (DAI)			
Slight / Defined	41 (59.4)	04 (57.1)	0.999*
Severe / Very severe	28 (40.6)	03 (42.9)	
Tooth extraction			
Yes	09 (13.0)	00 (00.0)	0.594**
No	60 (87.0)	10 (100.0)	
Outcome measures	Mean (SD)	Mean (SD)	
Overall B-IFAM (T1)	97.26 (24.41)	96.50 (24.11)	0.952**
Overall B-IFAM (T2)	98.28 (22.57)	-	-
Overall B-IFAM (change T1-T2)	-1.01 (18.11)	-	-

DAI=Dental Aesthetic Index; SD=standard deviation
*Fisher's test, **Student's *t* test

adolescents who participated in the study, 38 were female (55.1%) and 31 were male (44.9%). Mean age was 12.32 years (± 1.63). Fifty-five adolescents used a conventional fixed appliance, 13 used a self-ligating appliance, and one used an aesthetic device. Fig. 1 shows the flowchart of the study.




Fig. 1: Flowchart of the study

For the comparison between females and males, no significant difference was found for age, family monthly income, malocclusion severity, orthodontic tooth extraction, or the overall B-IFAM score in T1 ($p > 0.05$). For the overall B-IFAM score in T2 ($p = 0.003$), and for the difference of the overall B-IFAM score between T1 and T2 ($p = 0.039$), girls had a more negative perception of the OHRQoL than boys (Table 2).

In the adjusted regression model, girls had a significantly greater increase in overall B-IFAM score than boys, indicating a more negative perception of the OHRQoL during the observation time (T1 to T2) (Coefficient = 11.77, 95 % CI = 3.47 – 20.60, $p = 0.006$). In addition, a greater overall B-IFAM score in T1 was associated with a reduction in the overall B-IFAM score over time (coefficient = 15.62; 95 % CI = 7.55 – 23.69, $p = 0.001$) (Table 3). Comparison of the scores of the B-IFAM domains showed that females had a significantly greater increase in the score from T1 to T2 (more negative perception of the OHRQoL over time) than males for the aesthetic domain ($p = 0.034$) and physical impact domain ($p = 0.011$). As these were greater than the MCID, these differences were clinically significant. For these two domains, the effect size (the magnitude of the difference) was moderate (Table 4).

Table 2. Comparison of covariates and B-IFAM overall scores between female and male adolescents

	Girls	Boys	<i>p</i> value
Covariates	N (%)	N (%)	
Age			
≤12 years	26 (68.4)	18 (58.1)	0.453 [†]
>12 years	12 (31.6)	13 (41.9)	
Family Monthly income			
≤2 minimum wages	27 (71.1)	17 (54.8)	0.211 [†]
>2 minimum wages	11 (28.9)	1 (45.2)	
Malocclusion (DAI)			
Slight / Defined	25 (65.8)	16 (51.6)	0.325 [†]
Severe / Very severe	13 (34.2)	15 (48.4)	
Tooth extraction			
Yes	04 (10.5)	05 (16.1)	0.721 ^{**}
No	34 (89.5)	26 (83.9)	
Outcome measures	Mean (SD)	Mean (SD)	
Overall B-IFAM (T1)	99.09 (18.91)	93.54 (21.29)	0.241 ^{***}
Overall B-IFAM (T2)	103.31 (21.31)	89.77 (22.07)	0.003 ^{***}
Overall B-IFAM (change T1-T2)	-5.05 (18.10)	3.93 (17.12)	0.039 ^{***}

DAI=Dental Aesthetic Index; SD=standard deviation
[†]Pearson's test, ^{**}Fisher's test, ^{***}Student's *t* test

Table 3. Crude and adjusted associations between adolescents' sex and changes in the B-IFAM overall score

	Overall B-IFAM (change T1-T2)	Crude associations			Adjusted associations		
	Mean (SD)	Coefficient	95% CI	p value*	Coefficient	95% CI	p value*
Sex							
Girls	-5.05 (18.10)	Reference			Reference		
Boys	3.94 (17.12)	8.98	0.45 – 17.52	0.039	11.77	3.47 – 20.60	0.006
Age							
≤12 years	-1.27 (20.93)	Reference			Reference		
>12 years	-0.56 (11.97)	0.71	-8.40 – 9.83	0.876	-0.93	-9.76 – 7.90	0.834
Family Monthly income							
≤2 minimum wages	-1.61 (16.24)	Reference			Reference		
>2 minimum wages	0.04 (21.32)	1.65	-7.45 – 10.76	0.718	-1.79	-10.55 – 6.96	0.684
Malocclusion (DAI)							
Slight / Defined	-1.73 (16.39)	Reference			Reference		
Severe / Very severe	0.04 (20.63)	1.76	-7.15 – 10.68	0.694	-0.01	-8.26 – 8.25	0.999
Tooth extraction							
Yes	6.78 (18.41)	Reference			Reference		
No	-2.18 (17.92)	-8.96	-21.79 – 3.87	0.168	-10.76	-22.69 – 1.17	0.076
Overall B-IFAM (T1)							
Low impact (score ≤99)	-7.17 (14.41)	Reference			Reference		
High impact (score >99)	5.32 (19.49)	12.49	4.27 – 20.71	0.003	15.62	7.55 – 23.69	0.001

SD=standard deviation, CI=confidence interval

*Linear regression. Significant at $p < 0.05$ **Table 4. Comparison of the B-IFAM domain scores and the minimal clinically important difference between female and male adolescents**

	Number of questions (score range)	Girls Change T1-T2 Mean (SD)	Boys Change T1-T2 Mean (SD)	p value*	Difference between girls and boys Mean (95% CI)	Effect size 95% CI	MCID
Aesthetics	5 (5 – 25)	-0.63 (3.16)	1.00 (3.05)	0.034	-1.63 (-3.13 – -0.12)	0.50 (0.07 – 0.93)	1.60
Functional limitations	3 (3 – 15)	0.13 (2.50)	0.35 (2.87)	0.731	-0.22 (-1.51 – 1.06)	0.08 (-0.35 – 0.51)	1.32
Dietary impact	6 (6 – 30)	0.26 (5.36)	0.77 (5.97)	0.710	-0.51 (-3.23 – 2.21)	0.09 (-0.34 – 0.52)	2.80
Oral hygiene impact	3 (3 – 15)	-0.39 (3.14)	0.26 (3.36)	0.409	-0.65 (-2.21 – 0.91)	0.20 (-0.23 – 0.63)	1.61
Maintenance	2 (2 – 10)	-0.08 (2.04)	-0.01 (2.03)	0.873	-0.07 (-1.06 – 0.90)	0.03 (-0.40 – 0.46)	1.01
Physical impact	9 (9 – 45)	-2.89 (5.84)	0.94 (6.27)	0.011	-3.83 (-6.74 – -0.91)	0.60 (0.38 – 0.82)	3.14
Social impact	5 (5 – 25)	-0.42 (3.50)	0.58 (2.87)	0.205	-1.00 (-2.56 – 0.56)	0.30 (-0.13 – 0.73)	1.62
Time constraints	5 (5 – 25)	-1.08 (3.57)	0.09 (3.36)	0.167	-1.17 (-2.85 – 0.50)	0.33 (-0.10 – 0.76)	1.75
Travel/cost/ inconveniences	5 (5 – 25)	0.05 (4.20)	-0.06 (3.28)	0.900	0.11 (-1.72 – 1.96)	0.29 (-0.14 – 0.72)	1.89

SD=standard deviation, CI=confidence interval, MCID=minimally clinically important difference

*Student's *t* test. Significant at $p < 0.05$ **Table 5. Analysis of covariance comparing the B-IFAM domain scores between female and male adolescents, controlling for type of fixed appliance worn**

	Aesthetics	Functional limitations	Dietary impact	Oral hygiene impact	Maintenance	Physical impact	Social impact	Time constraints	Travel/cost/ inconve- niences
Sex									
Girls	-1.61 (0.75) *	-0.19 (0.65) *	-0.42 (1.36) *	-0.58 (0.77) *	-0.05 (0.49) *	-3.72 (1.45)	-0.97 (0.78) *	-1.16 (0.84) *	0.13 (0.93) *
Boys	0	0	0	0	0	0	0	0	0
p value	0.037	0.761	0.754	0.455	0.907	0.013	0.221	0.174	0.881
Type of fixed appliance									
Conventional	-0.39 (0.94) *	-0.59 (0.80) *	-1.95 (1.68) *	-1.70 (0.95) *	-0.49 (0.61) *	-2.57 (1.79) *	-0.70 (0.97) *	-0.16 (1.05) *	-0.54 (1.15) *
Self-lig/aesth	0	0	0	0	0	0	0	0	0
p value	0.674	0.464	0.251	0.080	0.419	0.156	0.469	0.873	0.637

Self-lig/aesth=self-ligating/aesthetic fixed appliance

*Coefficient (error)

p value=analysis of covariance. Significant at $p < 0.05$

The analysis of covariance showed that the increase in the score from T1 to T2 (more negative perception of the OHRQoL over time) in females than in males for the aesthetic domain ($p=0.037$) and physical impact domain ($p=0.013$) remained, regardless of the variable type of fixed appliance used (Table 5).

DISCUSSION

Significant changes were found in the OHRQoL of females over the first year of orthodontic treatment with a fixed appliance. After 12 months of wearing of the fixed appliance, girls had a more negative perception of the OHRQoL than boys. The literature shows favorable effects on the OHRQoL of adolescents, regardless of sex, after the first year of wearing a fixed appliance, showing improvement in their overall OHRQoL, as a result of the strong positive repercussions on their emotional and social wellbeing¹⁹. However, there are no longitudinal studies comparing the difference between sexes regarding the impact of the first year of wearing a fixed appliance on the OHRQoL. Studies that evaluated the impact of the need for orthodontic treatment and the OHRQoL of adolescents have shown that girls with occlusal disharmony have a more negative perception of their OHRQoL than boys in a similar condition. Females seem to have greater aesthetic concern and can feel more upset by dental disharmony and extremely exaggerated skeletal changes in the face²⁰. Although the literature shows that girls seek orthodontic treatment more often to correct malocclusion^{20,21}, the present study found that girls, when compared to boys, have a more negative perception of the OHRQoL between the third month and the first year of treatment with the fixed appliance. This finding illustrates that the more negative perception of the quality of life before orthodontic treatment^{20,22} seems to perpetuate itself throughout the 12 months after the fixed appliance has been installed. In the adjusted regression for the evaluated confounding variables, the girls continued to have a more negative perception of the OHRQoL throughout the orthodontic treatment with the fixed appliance, when compared to the boys. In this regard, a study conducted in 2014 evaluated the relationship between adolescents' OHRQoL and self-esteem during the first year of orthodontic treatment. It found that adolescents with better self-esteem at the beginning of the orthodontic treatment demonstrated minimal variability in the OHRQoL

after the first year of wearing a fixed appliance²³. The differences in self-esteem between adolescents of different sexes²⁴ may be useful to explain how orthodontic treatment impacts their quality of life²⁵. Females had a more negative perception of the OHRQoL during the first year of orthodontic treatment in the aesthetic domain. Adolescents undergoing orthodontic treatment with a fixed appliance seem to need aesthetic approval from their peers and parents. They often compare their looks to those of their friends and people on the media, and feel that their own teeth should be similar to the dental and facial aesthetic standard imposed by others²². The results of the present study suggest that girls feel more uncomfortable and embarrassed, and avoid smiling when wearing a fixed appliance, because they might perceive the orthodontic device as ugly.

Pain and discomfort in the oral mucosa are common complaints during orthodontic therapy. In the present study, girls reported more complaints than boys regarding pain and sores (physical impact) during the first year wearing a fixed appliance. A previous cross-sectional study using the B-IFAM questionnaire on patients in the sixth month of wearing a fixed appliance, also found differences in the perception of pain between boys and girls. It reported that girls appeared to be more anxious about the possibility of pain caused by the activation of the fixed appliance¹⁰. The results of the present study showed that this fear of pain caused by the activation of the fixed appliance persists in the girls even after the first year wearing it. It is important to highlight that the adolescents answer the questionnaire shortly after the dental appointment to activate the fixed appliance, which may influence the perception of pain, given that three days after activation, its perceived tension often subsides²⁶.

It is important to emphasize that the statistically significant differences between girls and boys for the domains of aesthetics and physical impact appear to be clinically significant, as they were greater than the MCID. The MCID reflects the changes of a specific outcome during a specific intervention, which are expressive and significant for the patients who undergo that type of intervention. In other words, it would be the smallest difference in a given outcome considered to be beneficial and which would imply, in the absence of side effects and excessive costs, a change in the management of the patient²⁷. In

studies that evaluate quality of life, a significant change in the OHRQoL is considered as one which results in a significant reduction in the symptoms or an improvement in function and wellbeing²⁸. This concept helps resolve one of the greatest challenges of transforming scientific evidence into practice, which is the interpretation of research data in the light of clinical relevance²⁹. Moreover, within our results, the MCID implications were confirmed by effect size for the two domains, which were moderate to high³⁰.

The results of studies on quality of life can be important to help the clinician to understand the physical, functional, and wellbeing consequences of orthodontic treatment. Based on the results of the present study, girls have a more negative perception of wearing a fixed appliance for reasons of aesthetics

and probable pain and discomfort caused by the orthodontic device. Thus, early advice regarding the repercussions of orthodontic treatment should place greater emphasis on girls, to provide greater understanding of the possible complications of wearing a fixed appliance¹⁰ to correct occlusal discrepancies, thereby improving the patient's quality of life at the end of the treatment³¹. Moreover, it is of utmost importance to clarify the effects of orthodontic therapy with a fixed appliance to the decision-makers, to enable improvement and better organization of the services providing orthodontic care⁹.

CONCLUSION

The impact of wearing a fixed appliance on OHRQoL was more negative in female than male adolescents during the first 12 months of orthodontic treatment.

ACKNOWLEDGMENTS

We thank the staff of the orthodontic clinic of the School of Dentistry of the Federal University of Minas Gerais.

FUNDING

Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Colegiado de Pós-Graduação em Odontologia (CPGO), and Pró-Reitoria de Pós-Graduação da Universidade Federal de Minas Gerais (PRPq/UFMG). S.M.P. and L.G.A. are research fellows of CNPq.

DECLARATION OF CONFLICTING INTERESTS

The authors declare no potential conflicts of interest regarding the research, authorship, and/or publication of this article

REFERENCES

1. Brogardh-Roth S, Paulsson L, Larsson P, Ekberg E. Do preterm-born adolescents have a poorer oral health-related quality of life? *BMC Oral Health*. 2021;21(1):440. <https://doi.org/10.1186/s12903-021-01799-3>
2. Haag DG, Peres KG, Balasubramanian M, Brennan DS. Oral condition and health-related quality of life: A systematic review. *J Dent Res*. 2017;96(8):864-74. <https://doi.org/10.1177/0022034517709737>
3. Yactayo-Albuquerque MT, Alen-Méndez ML, Azañedo D, Comandé D, Hernández-Vásquez A. Impact of oral diseases on oral health-related quality of life: A systematic review of studies conducted in Latin America and the Caribbean. *PLoS One*. 2021;16(6):e0252578. <https://doi.org/10.1371/journal.pone.0252578>
4. Alrashed M, Alqerban A. The relationship between malocclusion and oral health-related quality of life among adolescents: a systematic literature review and meta-analysis. *Eur J Orthod*. 2021;43(2):173-83. <https://doi.org/10.1093/ejo/cjaa051>
5. Sun L, Wong HM, McGrath CP. Relationship between the severity of malocclusion and oral health related quality of life: a systematic review and meta-analysis. *Oral Health Prev Dent*. 2017;15(6):503-17.
6. Davies SJ, Gray RM, Sandler PJ, O'Brien K. Orthodontics and occlusion. *Br Dent J*. 2001;91(10):545-91. <https://doi.org/10.1038/sj.bdj.4801229a>
7. Zhang M, Colman M, Hagg U. Changes in oral health-related quality of life during fixed orthodontic appliance therapy. *Am J Orthod Dentofacial Orthop*. 2008;133(1):23-9. <https://doi.org/10.1016/j.ajodo.2007.01.024>
8. Abreu LG, Lages EMB, Abreu MHNG, Pereira LJ, Paiva SM. Preadolescent's oral health-related quality of life during the first month of fixed orthodontic appliance therapy. *J Orthod*. 2013;40(3):218-24. <https://doi.org/10.1179/1465313313Y.0000000053>
9. Cunningham SJ, Hunt NP. Quality of life and its importance in orthodontics. *J Orthod*. 2001; 28(2):152-158. <https://doi.org/10.1093/ortho/28.2.152>
10. Corradi-Dias L, Paiva SM, Pretti H, Pordeus IA, Abreu LG. Impact of onset of fixed appliance therapy on adolescents' quality of life using a specific condition questionnaire: A cross-sectional comparison between male and female

- individuals. *J Orthod.* 2019;46(3):195-204. <https://doi.org/10.1177/1465312519851220>
11. Meade T, Dowswell E. Adolescents' health-related quality of life (HRQoL) changes over time: a three year longitudinal study. *Health Qual Life Outcomes.* 2016;14:14. <https://doi.org/10.1186/s12955-016-0415-9>
 12. Von EE, Altman DG, Egger M, Pocock SJ, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol.* 2008;61(4):344-49. <https://doi.org/10.1016/j.jclinepi.2007.11.008>
 13. Mandall NA, Vine S, Hulland R, Worthington HV. The impact of fixed orthodontic appliances on daily life. *Community Dent Health* 2006;23(2):69-74.
 14. Rebouças AP, Bendo CB, Abreu LG, Lages EMB, et al. Cross-cultural adaptation and validation of the impact of fixed appliances measure questionnaire in Brazil. *Brazil Oral Res.* 2018;32(14). <https://doi.org/10.1590/1807-3107bor-2018.vol32.0014>
 15. Jenny J, Cons NC. Establishing malocclusion severity levels on the Dental Aesthetic Index (DAI) scale. *Aust Dent J.* 1996;41(1):43-6. <https://doi.org/10.1111/j.1834-7819.1996.tb05654.x>
 16. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics* 1977;33(1):159-74. <https://doi.org/10.2307/2529310>
 17. Cohen J. *Statistical Power Analysis for the Behavioural Sciences.* Mahwah: 768 Lawrence Erlbaum Associates 1988.
 18. Copay AG, Subach BR, Glassman SD, Polly DW, et al. Understanding the minimum clinically important difference: a review of concepts and methods. *Spine J.* 2007;7(5):541-6. <https://doi.org/10.1016/j.spinee.2007.01.008>
 19. Abreu LG, Melgaço CA, Lages EMB, Abreu MHNG, et al. Effect of year one orthodontic treatment on the quality of life of adolescents, assessed by short form of Child Perceptions Questionnaire. *Eur Arch Paediatr Dent.* 2014;15:435-41. <https://doi.org/10.1007/s40368-014-0135-y>
 20. Feu D, Oliveira BH, Almeida MAO, Kiyak HA, et al. Oral health-related quality of life and orthodontic treatment seeking. *Am J Orthod Dentofacial Orthop.* 2010;138(2):152-9. <https://doi.org/10.1016/j.ajodo.2008.09.033>
 21. Alhaja ESJA, Al-Nimri KS, Al-Khateeb S. Self-perception of malocclusion among north Jordanian school children. *Eur J Orthod.* 2005;27(3):292-5. <https://doi.org/10.1093/ejo/cjh094>
 22. Ao H, Deng X, She Y, Wen X, et al. A biopsychosocial-cultural model for understanding oral-health-related quality of life among adolescents orthodontic patients. *Health Qual Life Outcomes.* 2020;18(1):86. <https://doi.org/10.1186/s12955-020-01334-y>
 23. Brosens V, Ghijeslings I, Lemiere J, Fieuws S, et al. Changes in oral health-related quality of life reports in children during orthodontic treatment and the possible role of self-esteem: a follow-up study. *Eur J Orthod.* 2014;36(2):186-91. <https://doi.org/10.1093/ejo/cjt035>
 24. Quatman T, Watson CM. Gender differences in adolescent self-esteem: an exploration of domains. *J Genet Psychol.* 2001;162(1):93-117. <https://doi.org/10.1080/00221320109597883>
 25. De Baets E, Lambrechts H, Lemiere J, Diya L, et al. Impact of self-esteem on the relationship between orthodontic treatment need and oral health-related quality of life in 11 - to 16-year-old children. *Eur J Orthod* 2012;34(6):731-7. <https://doi.org/10.1093/ejo/cjr088>
 26. Sergl HG, Klages U, Zentner A. Pain and discomfort during orthodontic treatment causative factors and effects on compliance. *Am J Orthod Dentofacial Orthop.* 1998; 114(6):684-91. [https://doi.org/10.1016/S0889-5406\(98\)70201-X](https://doi.org/10.1016/S0889-5406(98)70201-X)
 27. Jaeschke R, Singer J, Guyatt. Measurement of health status. Ascertain the minimal clinically important difference. *Control Clin Trials.* 1989;10(4):407-15. [https://doi.org/10.1016/0197-2456\(89\)90005-6](https://doi.org/10.1016/0197-2456(89)90005-6)
 28. Crosby RD, Kolotkin RL, Williams GR. Defining clinically meaningful change in health related quality of life. *J Clin Epidemiol.* 2003;56(5):395-407. [https://doi.org/10.1016/S0895-4356\(03\)00044-1](https://doi.org/10.1016/S0895-4356(03)00044-1)
 29. Rennard SI. Minimal clinically important difference, clinical perspective: an opinion. *COPD.* 2005; 2(1):51-5. <https://doi.org/10.1081/COPD-200050641>
 30. Kazis LE, Anderson JJ, Meenan RF. Effect sizes for interpreting changes in health status. *Med Care.* 1989;27(3 Suppl):S:178-89. <https://doi.org/10.1097/00005650-198903001-00015>
 31. Chen M, Wang DW, Wu LP. Fixed orthodontic appliance therapy and its impact on oral health-related quality of life in Chinese patients. *Angle Orthod.* 2010;80(1):49-56. <https://doi.org/10.2319/010509-9.1>