

# Modulation of IL-10/IL-10R expression by mafosfamide, a derivative of 4-hydroxycyclophosphamide, in a rat B-cell lymphoma

MARÍA J. RICO\*<sup>1,2</sup>, PABLO MATAR\*<sup>1,2</sup>, O. GRACIELA SCHAROVSKY<sup>1,3</sup>

1. Instituto de Genética Experimental, Facultad de Ciencias Médicas, Universidad Nacional de Rosario, Santa Fe 3100, (2000) Rosario, Argentina.
2. Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).
3. Consejo de Investigaciones, UNR; Rosario, Argentina.

\* Both authors contributed equally to this work

**Key words:** metastasis, cyclophosphamide, cell proliferation

**ABSTRACT:** We have already shown that IL-10 plays an important role in immunosuppression and metastatic dissemination in the rat B-cell lymphoma L-TACB model. It was suggested that the up-regulation of IL-10 production and IL-10 receptor (IL-10R) expression would be part of the transition from primary tumor to metastatic phenotype and that IL-10, besides its immunosuppressive activity, may act as a growth factor for metastatic L-TACB cells. The treatment of L-TACB-bearing rats with a single low-dose cyclophosphamide decreased IL-10 production, reverted immunosuppression and induced the immunologic rejection of tumor metastasis without any effect on primary tumor growth. Our current aim was to investigate the effects of cyclophosphamide on the expression of IL-10 and IL-10R on primary and metastatic L-TACB cells. Considering that cyclophosphamide is a prodrug, we used mafosfamide, a compound that yields *in vitro* the same active metabolites as cyclophosphamide does *in vivo*. Mafosfamide induced down-regulation of IL-10 production and IL-10R expression on metastatic cells and, concomitantly, inhibited metastatic cell proliferation. We suggest that mafosfamide would inhibit the regulatory loop mediated by the IL-10/IL-10R system and, as a consequence, metastatic cell proliferation. These results may have a considerable impact on the design of new therapies for metastatic lymphomas.

## Introduction

During tumor progression, cancer cells change (enhance or decrease) the expression of several growth factors, cytokines, and their specific receptors, as well as other soluble factors or cell membrane receptors, in

order to grow autonomously and to acquire competence to metastasis (Pals *et al.*, 2007). Interleukin-10 (IL-10) and its receptor (IL-10R) have been implicated in progression and metastasis of different tumor types, including lymphoma. It was shown that the molecular signature of mantle cell lymphoma (MCL), a non-Hodgkin lymphoma, included overexpression of IL-10R, and that IL-10 was able to sustain tumor cell survival and proliferation (Visser *et al.*, 2000). On the other hand, overproduction of IL-10 is strongly associated with the immunosuppression frequently observed in tumor-bearing hosts and it could be respon-

Address correspondence to: O. G. Scharovsky.  
E-mail: [graciela.scharovsky@gmail.com](mailto:graciela.scharovsky@gmail.com)

Received: May 2, 2012. Revised version received: July 14, 2012.  
Accepted: July 17, 2012.

sible for the tumor escape from immune rejection (Salazar-Onfray *et al.*, 2007).

We have proposed that the IL-10/IL10R system represents a promising therapeutic target for malignant lymphoma (Croci *et al.*, 2007). In our current experiments we used the rat B-cell lymphoma L-TACB model, in which IL-10 appears to play a pivotal role in immunosuppression and metastatic dissemination (Rico *et al.*, 2005), and its inhibition or activity modulation constitutes an important therapeutic option (Matar *et al.*, 2002). Our previous results in this lymphoma model showed a differential effect of cyclophosphamide on primary tumor or metastasis growth: a single low-dose cyclophosphamide inhibited metastasis development without any effect on primary tumor growth (Matar *et al.*, 1995). This antimetastatic action of cyclophosphamide was mediated by immunomodulation since this effect could be adoptively transferred by immune cells of cyclophosphamide-treated tumor-bearing rats and the same treatment did not induce any effect in immunodeficient nude mice (Matar *et al.*, 1998). We showed that IL-10 was the main factor responsible for the induction and development of immunosuppression in L-TACB-bearing hosts and that the antimetastatic immunomodulatory effect of cyclophosphamide was mediated by a reduction in IL-10 levels produced by T-lymphocytes (Matar *et al.*, 2000; Matar *et al.*, 2001). In fact, cyclophosphamide induced a Th2/Th1 switch and increased the proliferation rate of spleen cells (Matar *et al.*, 2002). Also, we have shown that L-TACB cells from metastatic nodes produced high amounts of IL-10 and expressed higher IL-10 receptor (IL-10R) levels than L-TACB cells from primary tumors, which did not produce IL-10 at all (Rico *et al.*, 2005). Also, the addition of IL-10 *in vitro* induced an increase in proliferation rate only on metastatic cells (Matar *et al.*, 2002). It was suggested that the up-regulation of IL-10 production and IL-10R expression would be part of the transition from primary tumor to metastatic phenotype and that IL-10, besides its immunosuppressive activity, may act as a growth factor for metastatic L-TACB cells. Altogether, these results suggested that, in addition to its immunomodulatory activity, cyclophosphamide may also be able to regulate the secretion and/or activity of IL-10 produced by metastatic cells.

Thus, our current aim was to investigate the *in vitro* effects of active metabolites of cyclophosphamide on the expression of IL-10 and its specific receptor IL-10R on both primary tumor and metastatic L-TACB cells.

## Materials and Methods

### Drugs

Considering that cyclophosphamide is a prodrug, we used mafosfamide (Baxter Oncology GmbH, Germany), a compound that yields *in vitro* the same active metabolites as cyclophosphamide does *in vivo* (Niemeyer *et al.*, 1984). Solutions of 2.5, 5, 10 and 20  $\mu\text{M}$  mafosfamide were prepared in RPMI-1640 culture medium (Gibco BRL, Gaithersburg, MD, USA).

### Tumor


L-TACB is a poorly differentiated B-cell lymphoma that arose spontaneously in an inbred *e* rat (Calderari *et al.*, 1991). It is maintained by serial subcutaneous grafting by trocar of 1 mm<sup>3</sup> tumor fragments (approximately 10<sup>6</sup> cells) in syngeneic rats. When L-TACB is injected s.c., metastases occur exclusively in regional (axillary or inguinal) homolateral or contralateral lymph nodes in 20–30% of the animals.

### Preparation of tumor single-cell suspensions

To obtain single-cell suspensions from primary tumor and metastasis, *e* rats were implanted subcutaneously with L-TACB and 21 days later primary tumors and lymph node metastases were excised and cell suspensions were obtained by mechanical disruption in RPMI-1640 culture medium. At this time of tumor progression, hematoxylin–eosin staining showed that normal lymphoid tissue in an organized form was completely replaced by tumor tissue in lymph nodes, and that the only non lymphoid cells appearing could be assigned to nervous tissue and blood vessels (data not shown).

### Cell proliferation

Primary and metastatic L-TACB cells were incubated at  $1 \times 10^4$  per well in 100  $\mu\text{l}$  RPMI-1640 culture medium with 10% fetal calf serum (FCS) (Natocor, Córdoba, Argentina) containing 0, 2.5, 5, 10 or 20  $\mu\text{M}$  mafosfamide, at 37°C and 5% CO<sub>2</sub> for 48 h in 96-well plates. Cell proliferation was determined by the colorimetric WST-1 assay (Roche Diagnostics, Mannheim, Germany) following manufacturer instructions. Determinations were done in quadruplicate.



**Figure 1.** Effect of mafosfamide on cell proliferation, IL-10 production and IL-10R expression of primary tumor cells (1A,1C,1E, respectively) and metastatic cells (1B,1D,1F) of B-cell lymphoma L-TACB. ANOVA: (B) mafosfamide 0 µM vs 5, 10 or 20 µM,  $P < 0.05$ ; (D) mafosfamide 0 µM vs 2.5, 5, 10 or 20 µM,  $P < 0.05$ ; (F) mafosfamide 0 µM vs 5, 10 or 20 µM,  $P < 0.05$ .

### Quantification of IL-10

The concentration of IL-10 was measured in conditioned media from primary tumor or metastatic L-TACB cells by a sandwich ELISA assay. Five x 10<sup>5</sup> cells of each cell type suspended in 5 ml of RPMI-1640 culture medium supplemented with 10% FCS and mafosfamide (0-20 μM) were seeded in 25 cm<sup>2</sup> culture flasks. After 48 h incubation at 37°C and 5% CO<sub>2</sub>, culture supernatants were collected and centrifuged. The cell-free conditioned media were aliquoted and stored at -20°C until used. The ELISA sandwich technique, that uses two different antibodies against different epitopes of IL-10, was carried out according to the instructions provided in the kit (OptEIA™ Rat IL-10 Set; Pharmingen, San Diego, CA, USA).

### Expression of IL-10 receptor (IL-10R)

The effect of mafosfamide on IL-10R expression on primary tumor or metastatic cells was determined by CELISA (cellular ELISA) assay. Cells were pre-incubated with 0, 2.5, 5, 10 or 20 μM mafosfamide during 48 h and then fixed at 5 x 10<sup>5</sup> cells/well onto 96 wells microtiter plates with 0.5% formaldehyde-buffer for 18 h at 37°C. Then, recombinant IL-10 (rIL-10) at a saturating concentration (5000 pg/ml) was added. The plates were incubated 2 h at room temperature, washed, blocked and then incubated for 1 h with anti-IL-10-biotin/streptavidin-peroxidase conjugate. After the addition of tetramethylbenzidine and H<sub>2</sub>O<sub>2</sub> and 30 min incubation at room temperature in the dark, the reaction was stopped with H<sub>2</sub>SO<sub>4</sub>. The optical density was determined at 450-570 nm. All samples were assayed in quadruplicate.

### Statistical analysis

ANOVA was performed using GraphPad Prism® version 3.03 (GraphPad Software, San Diego, CA, USA). Differences between groups were considered significant when P < 0.05.

## Results

### Effect of mafosfamide on primary and metastatic L-TACB cell proliferation

Mafosfamide did not show any effect on primary tumor cells proliferation (Fig. 1A). In contrast, meta-

static L-TACB cells were sensitive to the antiproliferative effect of mafosfamide in a dose-dependent manner, from 5 to 20 μM, when compared to cells incubated without mafosfamide (P<0.05) (Fig. 1B).

### Effect of mafosfamide on IL-10 production

Quantification of IL-10 showed non-detectable levels in conditioned media from primary tumor cells (Fig. 1C), while the IL-10 levels in conditioned media from metastatic cells were higher in control cells than in cells treated with different doses of mafosfamide (P<0.05) (Fig. 1D). Considering that mafosfamide exerts a dose-dependent antiproliferative effect on metastatic cells (Fig. 1B), IL-10 levels were normalized to cell number for each mafosfamide concentration. In this condition, we also observed a mafosfamide-induced dose-dependent reduction of IL-10/cell (data not shown).

### Effect of mafosfamide on IL-10R expression

The results showed that primary tumor cells expressed low levels of IL-10R, which were not significantly modified by either dose of mafosfamide (Fig. 1E). On the contrary, metastatic L-TACB cells expressed higher levels of IL-10R with respect to primary tumor cells (P<0.05). Moreover, incubation with mafosfamide decreased IL-10R expression on metastatic L-TACB cells at doses of 5 to 20 μM (P<0.05) (Fig. 1F).

## Discussion

In this rat lymphoma model, as well as in human lymphomas, IL-10 plays a central role in the inhibition of the antitumor immune response as well as a growth factor for metastasis. Nowadays, high levels of IL-10 are considered as significant unfavorable prognostic factor and an International Prognostic Score-independent risk factor for treatment failure in advanced Hodgkin lymphoma (Matar *et al.*, 2001; Rautert *et al.*, 2008). We have shown that the treatment of lymphoma-bearing rats with a single low-dose cyclophosphamide inhibited metastasis development, and that this effect was mediated by cyclophosphamide-induced immunomodulation, mainly through a reduction in IL-10 levels (Matar *et al.*, 1998; 2000). We have also indicated that IL-10 may act as a growth factor for metastatic tumor cells in addition to its immunosuppressive activity (Rico *et al.*, 2005). Similar results were obtained in leukemic mantle cell lymphoma (Visser *et al.*, 2000), B16 mouse-

melanoma model (García-Hernández *et al.*, 2002), and freshly resected primary tumor and metastases from melanoma patients (Yue *et al.*, 1997). As far as we know, we demonstrated here a novel effect of mafosfamide on the expression of IL-10 and its specific receptor on lymphoma cells. We observed that mafosfamide, a derivative of 4-hydroxycyclophosphamide, induced down-regulation of IL-10 production and IL-10R expression on metastatic tumor cells and, concomitantly, inhibited metastatic cell proliferation. Taking into account that IL-10 induces *in vitro* metastatic cell proliferation (Matar *et al.*, 2002), and that metastatic cells simultaneously express IL-10R and produce IL-10, an autocrine and/or paracrine loop would take place, thus stimulating, either directly or indirectly, metastatic cell proliferation. We suggest that mafosfamide would inhibit the regulatory loop mediated by the IL-10/IL-10R system and, as a consequence, it would inhibit metastatic cell proliferation. Considering that mafosfamide yields the same active metabolites as cyclophosphamide in equimolar concentrations, it can be hypothesized that cyclophosphamide *in vivo* would produce a similar effect than mafosfamide *in vitro*. Then, the effects of IL-10 we have previously observed in L-TACB-bearing rats, namely, immunosuppression and induction of metastatic cells proliferation, could have been abrogated when animals were treated with antimetastatic low doses of cyclophosphamide. These results may have considerable impact on development of new therapies for metastatic lymphomas in view that a single low-dose of cyclophosphamide, a treatment devoid of toxicity, would act inhibiting at least two mechanisms contributing to malignant growth and progression: escape to immune rejection and autonomous growth of metastatic tumor cells.

### Acknowledgements

This work was supported by a grant from Alberto J. Roemmers Foundation to Pablo Matar.

### References

- Calderari S, Font MT, Garrocoq O, Martinez S, Morini JC, Puche R, Tarrés MC (1991). The inbred IIM/Fm stock. *Rat News Letter* **25**: 28–29.
- Croci DO, Zacarías Fluck MF, Rico MJ, Matar P, Rabinovich GA, Scharovsky OG (2007). Dynamic cross-talk between tumor and immune cells in orchestrating the immunosuppressive network at the tumor microenvironment. *Cancer Immunology Immunotherapy* **56**: 1687–1700.
- García-Hernández ML, Hernández-Pando R, Gariglio P, Berumen J (2002). Interleukin-10 promotes B16-melanoma growth by inhibition of macrophage functions and induction of tumour and vascular cell proliferation. *Immunology* **105**: 231–243.
- Matar P, Celoria GC, Font MT, Scharovsky OG (1995). Antimetastatic effect of a single-low dose of cyclophosphamide on a rat lymphoma. *Journal of Experimental and Clinical Cancer Research* **14**: 59–63.
- Matar P, Rozados VR, Roggero EA, Bonfil RD, Scharovsky OG (1998). Modulation of the antimetastatic effect of a single low dose of cyclophosphamide on rat lymphoma. *Tumor Biology* **19**: 69–76.
- Matar P, Rozados VR, González AD, Dlugovitzky DA, Bonfil RD, Scharovsky OG (2000). Mechanism of antimetastatic immunopotentiality by low-dose cyclophosphamide. *European Journal of Cancer* **36**: 1060–1066.
- Matar P, Rozados VR, Gervasoni SI, Scharovsky OG (2001). Down regulation of T-cell-derived IL-10 production by low-dose cyclophosphamide treatment in tumor-bearing rats restores *in vitro* normal lymphoproliferative response. *International Immunopharmacology* **1**: 307–319.
- Matar P, Rozados VR, Gervasoni SI, Scharovsky OG (2002). Th2/Th1 switch induced by a single low-dose of cyclophosphamide in a rat metastatic lymphoma model. *Cancer Immunology Immunotherapy* **50**: 588–596.
- Niemeyer U, Engel J, Scheffler G, Molge K, Sauerbier D, Weigert W (1984). Chemical characterization of ASTA Z 7557 (INN mafosfamide, CIS-4-sulfoethylthio-cyclophosphamide), a stable derivative of 4-hydroxy-cyclophosphamide. *Investigational New Drugs* **2**: 133–139.
- Pals ST, de Gorter DJ, Spaargaren M (2007). Lymphoma dissemination: the other face of lymphocyte homing. *Blood* **110**: 3102–3111.
- Rauter R, Schinköthe T, Franklin J, Weihrauch M, Boll B, Pogge E, Bredenfeld H, Engert A, Diehl V, Re D (2008). Elevated pretreatment interleukin-10 serum level is an International Prognostic Score (IPS)-independent risk factor for early treatment failure in advanced stage Hodgkin lymphoma. *Leukemia & Lymphoma* **49**: 2091–2098.
- Rico MJ, Matar P, Bonfil RD, Calcaterra N, Scharovsky OG (2005). The transition to the metastatic phenotype of rat lymphoma cells involves up-regulation of IL-10 receptor expression and IL-10 secretion. *Clinical & Experimental Metastasis* **22**: 127–135.
- Salazar-Onfray F, López MN, Mendoza-Naranjo A (2007). Paradoxical effects of cytokines in tumor immune surveillance and tumor immune escape. *Cytokine & Growth Factor Reviews* **18**: 171–182.
- Visser HP, Tewis M, Willemze R, Kluijn-Nelemans JC (2000). Mantle cell lymphoma proliferates upon IL-10 in the CD40 system. *Leukemia* **14**: 1483–1489.
- Yue FY, Dummer R, Geertsen R, Hofbauer G, Laine E, Manolio S, Burg G (1997). Interleukin-10 is a growth factor for human melanoma cells and down-regulates HLA class-I, HLA class-II and ICAM-I molecules. *International Journal of Cancer* **71**: 630–637.

