

## *Physocephala inhabilis* (Diptera: Conopidae) as a parasitoid of *Megachile (Sayapis) bomplandensis* (Hymenoptera: Megachilidae) in Argentina

ALMADA, Valentín<sup>1</sup>, DEMARCHI, Lucrecia<sup>1</sup>, FERRERAS, Esteban O.<sup>1</sup>, STUKE, Jens-Hermann<sup>2</sup>, CLEMENTS, David K.<sup>3</sup> & LUCIA, Mariano<sup>1,4,\*</sup>

<sup>1</sup> División Entomología, Museo de La Plata, Universidad Nacional de La Plata, Edificio Anexo Museo, Unidades de Investigación FCNyM. La Plata, Argentina. \*E-mail: mlucia@fcnym.unlp.edu.ar

<sup>2</sup> Roter Weg 22, D-26789 Leer, Germany.

<sup>3</sup> 7 Vista Rise, Llandaff, Cardiff, CF5 2SD, U.K.

<sup>4</sup> CONICET, Consejo Nacional de Investigaciones Científicas y Técnicas. Argentina.

Received 30 - VI - 2020 | Accepted 24 - VIII - 2020 | Published 28 - IX - 2020

<https://doi.org/10.25085/rsea.790307>

### ***Physocephala inhabilis* (Diptera: Conopidae) como parasitoide de *Megachile (Sayapis) bomplandensis* (Hymenoptera: Megachilidae) en Argentina**

**RESUMEN.** El conópido *Physocephala inhabilis* (Walker) es registrado como parasitoide de la abeja nativa *Megachile (Sayapis) bomplandensis* (Durante). También se reporta el hallazgo de una especie sin identificar del género *Physocephala* parasitando a *Megachile (Chrysosarus)* sp. en Argentina.

**PALABRAS CLAVE.** Abeja silvestre. Anthophila. Nuevo registro de hospedador. Polinizador.

**ABSTRACT.** The conopid fly *Physocephala inhabilis* (Walker) is newly recorded as parasitoid of the native bee *Megachile (Sayapis) bomplandensis* (Durante). We also document an unidentified species of *Physocephala* parasitizing a *Megachile (Chrysosarus)* sp. in Argentina.

**KEYWORDS.** Anthophila. New host record. Pollinator. Wild bee.

The conopid flies are - as far as is known - solitary and internal parasitoids of insects, chiefly of aculeate wasps and bees (Hymenoptera). The females of these flies attack adult bees during flight and oviposit through the intersegmental membranes of the metasoma, where the parasitoid larvae will develop (*e.g.* De Meijere, 1904; Howell, 1967). There are currently some 129 recognized species of *Physocephala* Schiner around the world, of which 28 are present in the Neotropics and nine have been recorded in Argentina (Gibson et al., 2014). To date, 15 Neotropical species of this genus have been recorded parasitizing adults of various bee genera (Stuke et al., 2011; Stuke & Cardoso, 2013; Couto & Camillo, 2014; Plischuk et al., 2017; Stuke, 2017).

Bees of the family Megachilidae are pollinators of wildflowers as well as of several important crops worldwide. Some species are used commercially as managed crop pollinators (Pitts-Singer & Cane, 2011; Haider et al., 2014). The genus *Megachile* occurs in a wide diversity of habitats in all continents except Antarctica, ranging from lowland tropical rain forests and deserts to high elevation environments (Gonzalez et al., 2019). More than 430 species grouped in 31 subgenera are recorded for the Neotropical region (Raw, 2007; Gonzalez et al., 2018; Roig-Alsina, 2020). Most species of *Megachile* build their nests in pre-existing burrows or cavities using pieces of leaves that are used to create the breeding cells. However, members


of the subgenera *Chrysosarus* Mitchell and *Sayapis* Titus use other materials such as mud combined with leaves or petals or chewed leaf material with soil or small pebbles (Michener, 2007). Currently, eight species of *Megachile* have been cited as being parasitized by different species of *Physocephala* around the world (Stuke & Cardoso, 2013). In the Neotropics, the species *Megachile* (*Moureapis*) *maculata* Smith from Brazil has been recorded as host of *Physocephala inhabilis* (Walker) (Stuke & Cardoso, 2013) and of an unidentified conopid fly (Cardoso & Silveira, 2011). In this paper we add new records of *Physocephala* parasitizing two Argentinian species of *Megachile*.

As part of a study on the diversity of native bees associated with horticultural crops performed using trap-nests made of wooden blocks, on 4-III-2019 we found a dead female of *Megachile* (*Sayapis*) *bomplandensis* (Durante) inside a trap-nest of 8 mm in diameter (Fig. 2). On 27-X-2019, approximately nine months later, a single male of *P. inhabilis* emerged (Fig. 1). Also, on 17-XII-2013, we found a dead female of *Megachile* (*Chrysosarus*) sp. (Fig. 3) inside a trap nest of the same diameter. The adult of the conopid did not emerge but the puparium was morphologically similar to that of the *P. inhabilis* found inside the *M. bomplandensis* (Fig. 3). Thus, we assumed it was likely to be the same genus of conopid.

*Physocephala inhabilis* is widely distributed in the Neotropical region (Stuke, 2017) and in Argentina, where it has been recorded in the Mendoza, Tucumán and Río Negro provinces (Kröber, 1915 [as soror]; Camras & Parrillo, 1996; Gibson et al., 2014). Evidence suggests that *P. inhabilis*, a relatively small species of conopid, has a wide host range, parasitizing both medium-sized bees such as *Megachile* and *Centris* [i.e. *Megachile* (*Moureapis*) *maculata* Smith (Stuke & Cardoso, 2013), *M. (*Sayapis*) bomplandensis* (this paper), and *Centris analis* Fabricius (Santos et al., 2008; Couto & Camillo, 2014)] with body size-ranges from about 9-10 mm, as well as larger species such as *Centris vittata* (Fabricius) and *Epicharis bicolor* Smith (Santos et al., 2008), with body size ranges of 20-22 mm and 16-18 mm respectively. The time between the collection date of the dead bee and the emergence of the conopid fly in our study is similar to that recorded for other conopid species such as *Physocephala wulpi* Camras reared from *Xylocopa* (*Neoxylocopa*) *augusti* Lepeletier. *Physocephala wulpi* emerges in the spring from puparia which are collected during the late summer or early autumn, having overwintered within the dead host (Lucia et al., 2020).

#### Material examined

One ♂ of *Physocephala inhabilis* (Walker) ARGENTINA, La Plata, Buenos Aires, 34°54'39"S, 57°55'37"W 18 m.a.s.l., emerged on 27-X-2019, Col. M. Lucia, & V. Almada. (ex female of *Megachile* (*Sayapis*) *bomplandensis* collected on 4-III-2019); 1 Puparium of *Physocephala* spec., ARGENTINA, La Plata, Buenos Aires



**Fig. 1. *Physocephala inhabilis* male. a, habitus in lateral view; b, habitus in dorsal view; c, head and thorax in dorsal view; d, head in frontal view. Scale bars: (a-b) 2 mm; (c-d) 1 mm.**



**Fig. 2. Females of *Megachile* (*Sayapis*) *bomplandensis* (Durante). a, habitus in lateral view; b, head in frontal view; c, part of the metasoma containing puparium of *Physocephala inhabilis* in dorsal view. Scale bars: 2 mm.**

34°54'39"S, 57°55'37"W, 18 m.a.s.l., Col. M. Lucia (ex female of *Megachile* (*Chrysosarus*) sp. collected on 17-XII-2013). The specimen of *Physocephala* as well as the two adult females of *Megachile* and the conopid puparium, were deposited in the collection of División Entomología del Museo de La Plata, Argentina (MLP).



**Fig. 3. Female of *Megachile (Chrysosarus)* sp. a, habitus in lateral view; b, head in frontal view; c, part of the metasoma containing puparium of a *Physocephala* species in dorsal view. Scale bars: 2 mm.**

#### ACKNOWLEDGMENTS

We thank Victor Gonzalez and three anonymous reviewers for their insightful comments which improved the manuscript. Also, Unidad Vivero (FCNYM, UNLP) for their support. This work was partially supported by PICT 2016-1846. Financial support was provided to ML by the Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina (CONICET).

#### LITERATURE CITED

- Camras, S., & Parrillo, P.P. (1996) New *Stylogaster* and ranges of Conopidae (Diptera) from the Brazilian and Bolivian Amazonia. *Acta Amazonica*, **25**, 221-234.
- Cardoso, C.F., & Silveira, F.A. (2011) Nesting biology of two species of *Megachile (Moureapis)* (Hymenoptera: Megachilidae) in a semideciduous forest reserve in southeastern Brazil. *Apidologie*, **43**, 71-81.
- Couto, R.M., & Camillo, E. (2014) Deposições de óleo por fêmeas de *Centris analis* (Fabricius) (Hymenoptera: Apidae: Centridini) parasitadas por *Physocephala* spp. (Diptera: Conopidae). *EntomoBrasilis*, **7**, 81-85.
- De Meijere, J.C.H. (1904) Beiträge zur Kenntnis der Biologie und der systematischen Verwandtschaft der Conopiden. *Tijdschrift voor Entomologie*, **46**, 144-224.
- Gibson, J.F., Skevington, J.H., & Camras, S. (2014) Conopidae. *Biodiversidad de Artrópodos Argentinos Volumen 4*. (Roig-Juñent, S., Claps, L.E., & Morrone, J.J. eds.), pp. 491-497. Editorial INSUE-UNT, San Miguel de Tucumán, Argentina.
- Gonzalez, H.V., Griswold, T., & Engel, M.S. (2018) South American Leaf-cutter bees (Genus *Megachile*) of the subgenera *Rhyssemegachile* and *Zonomegachile*, with two new subgenera (Hymenoptera: Megachilidae). *Bulletin of the American Museum of Natural History*, **425**, 1-73.
- Gonzalez, H.V., Gustafson, G.T., & Engel, M.S. (2019) Morphological phylogeny of Megachilini and the evolution of leaf-cutter behavior in bees (Hymenoptera: Megachilidae). *Journal of Melittology*, **85**, 1-123.
- Haider, M., Dorn, S., Sedivy, C., & Müller, A. (2014) Phylogeny and floral hosts of a predominantly pollen generalist group of mason bees (Megachilidae: Osmiini). *Biological Journal of the Linnean Society*, **111**, 78-91.
- Howell, J.F. (1967) Biology of *Zodion obliquefasciatum* (Macq.) (Diptera: Conopidae): a parasite of the alkali bee, *Nomia melanderi* Ckll. (Hymenoptera: Halictidae). *Technical Bulletin (Washington Agricultural Experiment Station)*, **51**, 1-33.
- Kröber, O. (1915) Die indo-australischen und südamerikanischen *Physocephala*-Arten. *Archiv für Naturgeschichte, Abteilung A*, **81(4)**, 117-146.
- Lucia, M., Ramello, P.J., & Gonzalez, V.H. (2020) Brood development and nest parasitism of *Xylocopa (Neoxylocopa) augusti* Lepeletier (Hymenoptera: Apidae), a promising crop pollinator in Argentina. *Journal of Applied Entomology*, DOI: 10.1111/jen.12773.
- Michener, C.D. (2007) *The Bees of the World* (2nd ed.). Johns Hopkins University Press, Baltimore.
- Pitts-Singer, T., & Cane, J.H. (2011) The alfalfa leafcutting bee, *Megachile rotundata*: The world's most intensively managed solitary bee. *Annual Review of Entomology*, **56**, 221-237.
- Plischuk, S., Salvarrey, S., Arbulo, N., Santos, E., Skevington, J.H., Kelso, S., Revainera, P.D., Maggi, M.D., Invernizzi, C., & Lange, C.E. (2017) Pathogens, parasites, and parasitoids associated with bumble bees (*Bombus* spp.) from Uruguay. *Apidologie*, **48**, 298-310.
- Raw, A. (2007) An annotated Catalogue of the Leafcutter and Mason Bees (Genus *Megachile*) of the Neotropics. *Zootaxa*, **1601**, 1-127.
- Roig-Alsina, A.H. (2020) *Joergensenella*, a new subgenus of Neotropical *Megachile* (Hymenoptera: Megachilidae), with a key to Argentinean *Megachile* with specialized facial pollen-harvesting hairs. *Revista del Museo Argentino de Ciencias Naturales. n. s.*, **22(1)**, 21-46.
- Santos, A.M., Serrano, J.C., Couto, R.M., Rocha, L.S.G., Mello-Patiu, C.A., & Garófalo, C.A. (2008) Conopid Flies (Diptera: Conopidae) parasitizing *Centris (Heterocentris) analis* (Fabricius) (Hymenoptera: Apidae, Centridini). *Neotropical Entomology*, **37**, 606-608.
- Stuke, J.H. (2017) *World Catalogue of Insects. Volume 15. Conopidae (Diptera)*. E. J. Brill, Leiden, Boston.
- Stuke, J.H., & Cardoso, C.F. (2013) *Physocephala inhabilis* (Walker) (Diptera: Conopidae) parasitizing *Megachile (Moureapis) maculata* Smith (Hymenoptera: Megachilidae). *Studia Dipterologica*, **20**, 39-43.
- Stuke, J.H., Lucia, M., & Abrahamovich, A.H. (2011) Host records of *Physocephala wulpi* Camras, with a description of the puparium (Diptera: Conopidae). *Zootaxa*, **3038**, 61-67.