

First record of *Orsillus depressus* (Hemiptera: Lygaeidae: Orsillinae) from the New World and Southern Hemisphere

D'HERVÉ, Federico E.^{1,2}; FERNÁNDEZ, Celeste Giselle¹ & DELLAPÉ, Pablo M.^{3,*}

¹ Laboratorio Regional de Plagas, SENASA. Ruta Nacional N° 22 km 1127 Villa Regina (8336), Río Negro, Argentina.

² Cátedra de Zoología Agrícola, Facultad de Ciencias Agrarias, Universidad Nacional del Comahue. Ruta Nacional N° 151 km 12,5 Cinco Saltos (8303), Río Negro, Argentina.

³ División Entomología, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, CONICET, Paseo del Bosque s/n, B1900FWA, La Plata, Argentina. *E-mail: pdellape@fcnym.unlp.edu.ar

Received 11 - VI - 2022 | Accepted 20 - IX - 2022 | Published 30 - XII - 2022

<https://doi.org/10.25085/rsea.810408>

Primer registro de *Orsillus depressus* (Hemiptera: Lygaeidae: Orsillinae) de América y del Hemisferio Sur

RESUMEN. El ligeido *Orsillus depressus* Dallas se alimenta de semillas de varias especies, principalmente cupresáceas. Desde finales del siglo pasado ha extendido su área de distribución desde su área de origen en la cuenca mediterránea hacia el noreste de Europa, y se la considera una especie invasora. En el presente trabajo se reporta por primera vez la presencia de *O. depressus* en el Nuevo Mundo y en el Hemisferio Sur. Se brindan datos de sus plantas hospedadoras y su distribución desde el centro-sur de Argentina. Además, se aportan caracteres morfológicos que permiten su identificación.

PALABRAS CLAVE. Argentina. Chinchas. Cupressaceae. Especies invasoras.

ABSTRACT. The lygaeoid *Orsillus depressus* Dallas feeds on seeds of several species, mainly cypress. Since the end of the last century, it has extended its range from its native area in the Mediterranean Basin towards northeastern Europe and it is considered an invasive species. In the present work, the presence of *O. depressus* is reported for the first time from the New World and Southern Hemisphere. Host plants and distribution from central-southern Argentina are given. In addition, morphological characters that allow their identification are provided.

KEYWORDS. Argentina. Cupressaceae. Invasive species. True bug.

The genus *Orsillus* Dallas (Hemiptera: Lygaeidae: Orsillinae) includes six species of medium to small-sized true bugs distributed in the Palearctic Region (Péricart, 1998, 2001; Dellapé & Henry, 2022). The nymphs and adults live in the cones of different species of conifers and use their stylets to feed on the seeds, causing damage to seed orchards, natural stands, and plantations (Roques et al., 1999; Battisti et al., 2000). In addition, when adults migrate in search of food resources or egg laying sites, they can carry spores of the fungus *Seiridium cardinale* Sutton & Gibson and *S. unicorn* (Cooke & Ellis) B. Sutton that causes cypress bark canker (Roques & Battisti, 1999; Ramos & Abrantes, 2000).

Orsillus depressus Dallas is the most widely distributed species in the genus (Péricart, 2001; Costas et al., 2018; Dellapé & Henry, 2022). Since the end of the last century, it has extended its range from its native area in the Mediterranean Basin towards northeastern Europe (Aukema, 1988; Hawkins, 1989; Korcz, 2007; Hebda et al., 2016; Rabitsch, 2008; Lis & Stolarczyk, 2018), and its presence in the Italian Alps is an example of its ability to adapt to harsher climates (Dioli, 1991). *Orsillus depressus* is considered a pest of conifers and an invasive species in Europe (Sweet, 2000; Wittenberg, 2005; CABI, 2022), and is on the list of quarantine pests in the USA (USDA, 2017).

Orsillus depressus appears to be less host-specific than other *Orsillus* species, being capable of feeding and ovipositing on species of *Chamaecyparis* Spach, *Cupressus* L., *Juniperus* L., *Sequoiadendron giganteum* (Lindley) J. Buchholz, *Thuja* L. (Cupressaceae), and *Pinus* L. (Pinaceae), including species introduced into its native area of the Mediterranean Basin (Péricart, 1998; Sweet, 2000; Rouault et al., 2005; Rabitsch, 2008). In Argentina, host conifers of *O. depressus* were introduced from the Northern Hemisphere for ornamental purposes (Parodi & Dimitri, 1972), being frequent in most parks, gardens and recreational areas. Some of these exotic species, such as *Cupressus arizonica* Greene and *C. macrocarpa* Hartw. ex Gord., are considered invasive in Córdoba Province, whereas others, such as *Juniperus communis* L., develop adventitiously in the Andean forests, (Giorgis & Tecco, 2014; Cantero et al., 2016) where there are also native Cupressaceae forests comprising *Austrocedrus chilensis* (D. Don) Pic. Serm. & Bizzarri, *Pilgerodendron uviferum* (D. Don) Florin, and *Fitzroya cupressoides* (Molina) I.M. Johnst.

Females of *O. depressus* lay their eggs in two or three-year-old cones, when they begin to open to disperse their seeds, or in the emergence holes of *Megastigmus wachtli* Seitner (Megastigmidae), a wasp whose larvae develop inside the cones (Roques & Battisti, 1999; Rouault et al., 2005). Another hymenopteran associated with *O. depressus* is the egg parasitoid *Telenomus* gr. *floridanus* (Scelionidae) (Rouault et al., 2000).

A female collected by the senior author on May 8, 2020, in Villa Regina town, Río Negro, Argentina, and identified as *O. depressus* triggered the search for more specimens. Between November 2021 and March 2022, samples were taken from trees in the family Cupressaceae to verify the presence of eggs, nymphs, and adults.

Surveys were carried out at 25 sites located in urban and suburban areas within a distance of 302 km in a north-south direction and 652 km in an east-west direction, covering 15 localities belonging to Buenos Aires, La Pampa, Neuquén, and Río Negro provinces. These localities are in most cases separated by extensive areas of grasslands and xerophytic shrubby plains.

Host trees belonging to the genera *Chamaecyparis*, *Cupressus*, and *Thuja* were identified at each site and 30 cones were collected, selecting those with open scales and still containing their seeds. On some occasions, samples were collected on different dates and hosts for the same site. In the laboratory, the samples from each site/date/host were placed in 1-liter plastic containers with cloth lids and observed with a Zeiss Stemi 2000-C 6.5x-50x stereomicroscope. Additionally, the cones were observed externally for holes that indicate the presence of species that provide oviposition sites, such as *M. wachtli*. Those cones that contained masses of unhatched eggs were kept in the laboratory at 25 °C for 30 days under 16L:8D conditions to observe the emergence of oophagous parasitoids. The collection sites were georeferenced by GPS device and the data obtained were

used to make a distribution map using the QGIS software version 3.8.1. The presence of *O. depressus* was communicated to the Argentine phytosanitary authorities through identification N° 580 of Sistema Nacional de Vigilancia y Monitoreo (SINAVIMO) and the examined specimens were deposited in the Museo de La Plata (MLP) La Plata, Buenos Aires, Argentina and in Laboratorio Regional de Plagas of Servicio Nacional de Sanidad Agroalimentaria (SENASA) in Villa Regina, Río Negro, Argentina.

Following Péricart's key to adults in the genus (1998), specimens were identified as *O. depressus* (Fig. 1). The first couplet in the key refers to "Rostrum not exceeding or slightly exceeding the middle of the abdomen; sternites III and IV with a more or less distinct longitudinal median groove." to *O. depressus* or *O. reyi* Puton or "Rostrum reaching or nearly apex of abdomen; sternites III and IV without a longitudinal median groove; second antennal segment of the order of 1.5 times as long as fourth" to *O. maculatus* (Fieber) and *O. pinicanariensis* Lindberg.



Fig. 1. *Orsillus depressus* female.

Our specimens show variability on rostral length, ranging from attaining the base of sternite VI to almost attaining the pygophore in some males. Rouault et al. (2005) noted that in areas where *C. sempervirens* occurs, *O. depressus* and *O. maculatus* may occur on the same tree, but morphological variability of adults and nymphs often hinders accurate identification of the species. They

measured the body length, rostral length, as well as their ratio (R:Bd) from several hundred specimens on different host plants, and found that the ratio was significantly lower in *O. depressus* (mean R:Bd < 0.85, although the values ranged from 0.62 to 0.89) than in *O. maculatus* (mean R:Bd > 0.85 in females and most males). Therefore, they argued that based on frequency distribution, this ratio does not seem to necessarily result in an accurate identification, particularly of males, and it is related to some degree with the host and the geographical location. According to these authors, the R:Bd ratio on fourth- and fifth-instar nymphs also could represent a diagnostic character for specific identification.

Besides the variability exhibited in rostral length and minor variation on coloration pattern, we believe that all specimens belong to the same specific unit because of their consistent morphology and vestiture. Most of the specimens lack a well-demarcated dark medial line on the pronotum, often used as diagnostic to identify *O. depressus*, and the dark area is more or less confined between the calli in most specimens, as described for *O. maculatus*, or extending somewhat diffusely over the posterior pronotal lobe. All specimens, however, show a more or less distinct longitudinal median groove on sternites III and IV, the pedicel is similar in length to the distiflagellomere and, the ratio body length-rostral length agrees to those found by Rouault et al. (2005) for *O. depressus*. We measured 20 males and 20 females from different localities and host, and we found a ratio from 0.72 to 0.90 in males (with the higher values from males collected on *C. sempervirens*), and a ratio from 0.67 to 0.83 in females. Additionally, we measured ten fifth-instar nymphs whose R:Bd ratio varied between 0.67 and 0.93,

within the set of measurements recorded for *O. depressus* (Rouault et al., 2005: Fig. 7)

The species is recorded from Buenos Aires, La Pampa, Neuquén, and Río Negro provinces in Argentina (Fig. 2). These records represent the first occurrence of this genus and species in the New World and Southern Hemisphere.




Fig. 2. Geographical distribution of *Orsillus depressus*: sites where was collected (red dots). Top left, area of origin in the Mediterranean basin and new distribution area in South America.

Of the 25 sites sampled, *O. depressus* was found in 18 (72%), which reveals its wide distribution in the study area (Table I). In addition, immature stages were observed in all these sites, indicating that this species is actively reproducing. No emergence holes or adults of the hymenopteran *M. wachtl* were found in the observed cones, nor were parasitoids obtained from the eggs.

Table I. Sites sampled in each province. Indicated: location, date (dd/mm/yyyy), geographic coordinates, altitude (meters above sea level), number of males (♂), females (♀), nymphs (n), egg masses (e) present in each sample and host species.

Site	Location	Date	Coordinates	Altitude	Individuals	Host
Buenos Aires						
1	Monte Hermoso	21/11/2021	38°59' 13'' S 61°17' 21'' W	15	0	<i>Cupressus sempervirens</i>
2	Monte Hermoso	21/11/2021	38°59' 04'' S 61°17' 22'' W	9	2♂ 5♀ 7n 2e	<i>Cupressus sempervirens</i>
3	Bahía Blanca	26/02/2022	38°43' 22'' S 62°15' 12'' W	23	1♂ 0♀ 58n 5e	<i>Cupressus sempervirens</i>
4	Pehuen C6	28/02/2022	38°59' 49'' S 61°35' 37'' W	17	0	<i>Cupressus sempervirens</i>
La Pampa						
5	Casa de Piedra	16/11/2021	38°09' 20'' S 67°08' 58'' W	294	6♂ 2♀ 95n 16e	<i>Cupressus lusitanica</i>
6	Gdor. Duval	11/02/2022	38°44' 40'' S 66°26' 11'' W	208	4♂ 1♀ 1n 3e	<i>Cupressus sempervirens</i>
7	Gdor. Duval	11/02/2022	38°44' 34'' S 66°26' 00'' W	207	1♂ 0♀ 9n 6e	<i>Cupressus arizonica</i>
8	Jacinto Arauz	01/03/2022	38°03' 18'' S 63°25' 31'' W	171	19♂ 10♀ 124n 35e	<i>Cupressus sempervirens</i>
Neuquén						
9	Neuquén	22/01/2022	38°57' 04'' S 68°02' 10'' W	272	0	<i>Cupressus arizonica</i>
10	Neuquén	22/01/2022	38°56' 51'' S 68°02' 46'' W	285	0	<i>Cupressus sempervirens</i>
11	Neuquén	22/01/2022	38°56' 30'' S 68°03' 32'' W	311	0	<i>Cupressus arizonica</i>
12	Neuquén	22/01/2022	38°56' 59'' S 68°03' 32'' W	285	1♂ 0♀ 12n 2e	<i>Cupressus arizonica</i>
13	Neuquén	22/01/2022	38°54' 22'' S 68°04' 58'' W	316	0♂ 1♀ 10n 4e	<i>Cupressus arizonica</i>
14	El Choc6n	05/03/2022	39°14' 48'' S 68°47' 56'' W	449	2♂ 0♀ 2n 1e	<i>Cupressus sempervirens</i>
15	El Choc6n	05/03/2022	39°15' 37'' S 68°49' 42'' W	437	1♂ 0♀ 2n 3e	<i>Cupressus sempervirens</i>

Table 1 (cont.). Sites sampled in each province. Indicated: location, date (dd/mm/yyyy), geographic coordinates, altitude (meters above sea level), number of males (♂), females (♀), nymphs (n), egg masses (e) present in each sample and host species.

Site	Location	Date	Coordinates	Altitude	Individuals	Host
Río Negro						
16	Villa Regina	08/05/2020 9/02/2022	39°06'08''S 67°02'17''W	203	1 ♀ 13♂ 11♀ 32n 6e	N/D <i>Thuja orientalis</i>
17	Villa Regina	24/11/2021 11/02/2022	39°05'35''S 67°04'50''W	217	13♂ 11♀ 32n 6e 2♂ 1♀ 5n 2e	<i>Cupressus arizonica</i> <i>Cupressus sempervirens</i>
18	Villa Regina	24/11/2021	39°09'52''S 67°06'16''W	210	0	<i>Cupressus arizonica</i>
19	Villa Regina	24/11/2021	39°06'05''S 67°04'43''W	206	0	<i>Cupressus sempervirens</i>
20	Viedma	02/01/2022	40°46'22''S 63°14'55''W	6	2♂ 6♀ 2n 12e	<i>Cupressus sempervirens</i>
21	Choele Choel	23/01/2022	39°16'53''S 65°39'58''W	139	7♂ 1♀ 15n 10e	<i>Cupressus arizonica</i>
22	Chimpay	23/01/2022	39°09'41''S 66°08'35''W	160	2♂ 4♀ 8n 7e	<i>Cupressus arizonica</i>
23	Cipolletti	14/01/2022	38°56'01''S 68°00'08''W	268	7♂ 0♀ 8n 12e	<i>Cupressus sempervirens</i>
24	General Roca	30/01/2022	39°02'29''S 67°35'45''W	234	4♂ 4♀ 24n 14e 10♂ 5♀ 38n 22e	<i>Cupressus arizonica</i> <i>Chamaecyparis lawsoniana</i>
25	Río Colorado	26/02/2022	38°43'22''S 62°15'12''W	83	2♂ 4♀ 3n 3e 4♂ 3♀ 11n 2e	<i>Cupressus arizonica</i> <i>Cupressus sempervirens</i>

Unlike what was reported in Europe where *O. depressus* spreads through the native vegetation, the arrival and dispersion of this species in Argentina requires human assistance in commerce to extend its range. *Orsillus depressus* can be considered as beneficial when feeding on the seeds of invasive exotic Cupressaceae. However, its feeding habits and ability to act as a vector of cryptogamic diseases reverse its role and makes it a serious threat to native cupressaceous forests considered vulnerable or endangered by the IUCN (2022).

ACKNOWLEDGMENTS

We thank Thomas J. Henry (Systematic Entomology Laboratory, ARS, USDA c/o National Museum of Natural History (NMNH), Washington, DC), who kindly reviewed the manuscript and offered comments for its improvement.

LITERATURE CITED

- Aukema, B. (1988) *Orsillus depressus* new for the Netherlands and Belgium (Heteroptera: Lygaeidae). *Entomologische Berichten*, **48**, 12, 181-183.
- Battisti, A., Cantini, R., Feci, E., Frigimelica, G., Guido, M., & Roques, A. (2000) Detection and evaluation of seed damage of cypress, *Cupressus sempervirens* L., in Italy. *Seed Science and Technology*, **28**, 199-208.
- CABI, Centre for Agricultural Bioscience International. (2022) *Invasive Species Compendium*. <https://www.cabi.org/isc/datasheet/116010> (Accessed on 10 March 2022).
- Cantero, J.J., Barboza, G., Chiarini, F., Deanna, R., Ariza Espinar, L., Giorgis, M., Núñez, C., & Bernardello, G. (2016) Novedades para la Flora de la Argentina. *Boletín de la Sociedad Argentina de Botánica*, **51**, 1, 183-207.
- Costas, M., López, T. & Vázquez, M.A. (2018) Checklist of Iberian Fauna. Superfamily Lygaeoidea Schilling, 1829 (Insecta: Heteroptera) in the Iberian peninsula, Balearic Islands and Canary Islands. *Fauna Ibérica*, **7** (ed.

Ramos, M.A. & Sánchez Ruiz, M.), 29 pp. Museo Nacional de Ciencias Naturales, CSIC. Madrid.

Dellapé, P.M., & Henry, T.J. (2022) *Lygaeoidea Species File*. <http://Lygaeoidea.SpeciesFile.org> (Accessed on 10 March 2022).

Dioli, P. (1991) Presenza di *Orsillus depressus* Dallas, 1852 nella zona alpina e osservazioni sulle specie italiane del genere (Insecta, Heteroptera, Lygaeidae). *Il Naturalista Valtellinese*, **2**, 47-51.

Giorgis M.A., & Tecco, P.A. (2014) Árboles y arbustos invasores de la Provincia de Córdoba (Argentina): una contribución a la sistematización de bases de datos globales. *Boletín de la Sociedad Argentina de Botánica*, **49**, 581-603.

Hawkins, R.D. (1989) *Orsillus depressus* Dallas (Hem., Lygaeidae) an arboreal groundbug new to Britain. *Entomologist's Monthly Magazine*, **125**, 241-244.

Hebda, G., Gierlasiński, G. & Woski, A. (2016) *Orsillus depressus* (Mulsant & Rey, 1852) (Hemiptera: Heteroptera: Lygaeidae) - pierwsze stwierdzenie na Dolnym Śląsku. *Heteroptera Poloniae - Acta Faunistica*, **10**, 23-24.

IUCN (2022) *The IUCN Red List of Threatened Species*. Version 2021-3. <https://www.iucnredlist.org>. (Accessed 10 March 2022).

Korc, A. (2007) *Orsillus depressus* (Mulsant & Rey) (Hemiptera: Heteroptera: Lygaeidae)-gatunek śródziemnomorski nowy dla Polskiej fauny. XLVI Zjazd Polskiego Towarzystwa Entomologicznego, Poznań, 20-22 września, Streszczenia Posterów, s. 30.

Lis, B., & Stolarczyk, T. (2018) *Orsillus depressus* (Mulsant et Rey, 1852) (Hemiptera: Heteroptera: Lygaeidae) pierwsze stwierdzenie na Nizinie Mazowieckiej. *Heteroptera Poloniae - Acta Faunistica*, **12**, 81-82.

Parodi, J., & Dimitri J.M. (1972) Descripción de las plantas cultivadas Cap IV. *Enciclopedia Argentina de Agricultura y Jardinería Vol. 1* ACME, Buenos Aires, Argentina. pp 74-100.

- Péricart, J. (1998) Hémiptères Lygaeidae euro-méditerranéens. Volume 1. Généralités. Systématique: première partie. Faune de France. France et régions limitrophes. 84A. Fédération Française des Sociétés de Sciences Naturelles, Paris, France.
- Péricart, J. (2001) Lygaeidae. *Catalogue of the Heteroptera of the Palearctic Region. Volume 4. Pentatomomorpha I.* (eds. Aukema B., & Rieger C.), pp. 35-220. The Netherland Entomological Society, Amsterdam.
- Rabitsch, W. (2008) Alien true bugs of Europe (Insecta: Hemiptera: Heteroptera). *Zootaxa*, **1827**, 1-44.
- Ramos, P., & Abrantes, C. (2000) Transmissão de esporos de fungos do género *Seiridium* por *Orsillus* spp. (Heteroptera: Lygaeidae) em Portugal. *En: Comunicações do III Encontro Nacional de Plantas Ornamentais, 2000*, Viana do Castelo, pp. 235-244.
- Roques, A., & Battisti, A. (1999) Cypress insect pests. *Cypress. A practical handbook.* (ed. E. Teissier du Cros, Ducrey, D., Barthelemy, C., Pichot, R., Giannini, P., Raddi, A., Roques, J., Sales, L., & B. Thibaut), pp. 74-95. Studio Leonardo, Florence.
- Roques, A., Markalas, S., Roux, G., Pan, Y., Sun, J., & Raimbault, J. (1999) Impact of insects damaging seed cones of cypress, *Cupressus sempervirens*, in natural stands and plantations of southeastern Europe. *Annals of Forest Science*, **56**, 167-177.
- Rouault, G., Roversi, P.F., Cantini, R., Battisti, A., Bouaziz, K., & Roques, A. (2000) First record of an egg parasitoid (*Telenomus* gr. *floridanus*, Hymenoptera Scelionidae) of two true seed bugs (Heteroptera Lygaeidae) living on Cupressaceae. *Redia*, **83**, 163-173.
- Rouault, G., Cantini, R., Battisti, A., & Roques, A. (2005) Geographic distribution and ecology of two species of *Orsillus* (Hemiptera: Lygaeidae) associated with cones of native and introduced Cupressaceae in Europe and the Mediterranean Basin. *Canadian Entomologist*, **137**, 4, 450-470.
- Sweet II, M.H. (2000) Seed and chinch bugs (Lygaeoidea). *Heteroptera of economic importance.* (ed. Schaefer, C.W., & Panizzi, A.R.), pp. 143-264. CRC Press, Boca Ratón.
- USDA-APHIS (2017) U.S. *Regulated Plant Pest List* <https://www.aphis.usda.gov/aphis/ourfocus/planthealth/import-information/rppl/rppl-table> (Accessed on 10 March 2022).
- Wittenberg, R. (ed). (2005) An inventory of alien species and their threat to biodiversity and economy in Switzerland. CABI Bioscience Switzerland Centre report to the Swiss Agency for Environment, Forest and Landscape. The environment in practice no. 0629. Federal Office for the Environment, Bern.